

Transfer matrices for scalar fields on curved spaces

E. Prodan

University of Houston, 4800 Calhoun Rd, Houston TX 77204-5508, USA

(February 7, 2008)

arXiv:math-ph/9901003v2 20 Jan 1999

Typeset using REVTeX

Abstract

We apply Nelson's technique of constructing Euclidean fields to the case of classical scalar fields on curved spaces. It is shown how to construct a transfer matrix and, for a class of metrics, the basic spectral properties of its generator are investigated. An application concerning decoupling of non-convex disjoint region is given.

I. INTRODUCTION

We start our construction from the ideas comprised in Nelson's axioms [1] for scalar Euclidean-Markoff quantum fields. Here, the Markoff property of certain projectors is one of the basic ingredient in defining the transfer matrix of whom generator is identified with the Hamiltonian of Wightman quantum scalar field. We found that these ideas can be used in the same way at the non-quantum level. In the case of the scalar fields on Riemannian manifolds, for an arbitrary direction, we construct a propagator by using the Markoff property. In the stationary case it becomes a semigroup which can be considered as the transfer matrix of the system and, further, it can be used in introducing a Hamiltonian. We will show that the propagator is exponentially bounded by using Agmon's [2] results in exponential decay of solutions of second-order elliptic equations. An application concerning the decoupling (in the sense of [3]) of two disjoint non-convex regions is given.

II. INTRODUCTORY DEFINITIONS AND RESULTS

Let us consider the Riemannian manifold (R^{n+1}, g) and the Laplace-Beltrami operator on it, Δ . For a point in R^{n+1} we use the notation (t, x) . Let $E_m(t, x; s, y)$ be the kernel of $(\Delta + m^2)^{-1}$ on $L^2(R^{n+1}, \sqrt{g} dt dx)$. As in [4], we will not consider the additional term $\frac{1}{6}\rho$. One defines the space $N \subset \mathcal{D}'(R^{n+1})$, $f \in N$ if :

$$\|f\|_N^2 = \int_{R^{n+1}} \int_{R^{n+1}} \bar{f}(t, x) E_m(t, x; s, y) f(s, y) \sqrt{g(t, x)} \sqrt{g(s, y)} dt dx ds dy < \infty, \quad (1)$$

and, for each $\sigma \in R$, let $N_\sigma \subset D'(R^n)$ be the space: $g \in N_\sigma$ if

$$\|g\|_{N_\sigma}^2 = \int_{R^n} \bar{g}(x) E_m(\sigma, x; \sigma, y) g(y) \sqrt{g(\sigma, x)} \sqrt{g(\sigma, y)} dx dy < \infty. \quad (2)$$

We will consider that, as in the Euclidean case, the space $L^2(R^n, d\mu_\sigma) \subset N_\sigma$, where $d\mu_\sigma(x) = \sqrt{g(\sigma, x)} d^n x$ and that it is dense in N_σ for each $\sigma \in R$. Now, let $\hat{E}_\sigma : N_\sigma \rightarrow L^2(R^n, d\mu_\sigma)$ be the operator corresponding to the kernel $E_m(\sigma, x; \sigma, y)$. Then $\hat{E}_\sigma^{1/2}$ defines an isometry from N_σ to $L^2(R^n, d\mu_\sigma)$ and let $(\hat{E}_\sigma^{1/2})^\dagger : L^2(R^n, d\mu_\sigma) \rightarrow N_\sigma$ be its adjoint. The following are true:

$$\hat{E}_\sigma^{1/2} \circ (\hat{E}_\sigma^{1/2})^\dagger = 1_{L^2(R^n, d\mu_\sigma)} \text{ and } (\hat{E}_\sigma^{1/2})^\dagger \circ \hat{E}_\sigma^{1/2} = 1_{N_\sigma}. \quad (3)$$

With our assumptions, $\hat{E}_\sigma^{1/2}(N_\sigma) = L^2(R^n, d\mu_\sigma) \subset N_\sigma$, the operator $\hat{E}_\sigma^{1/2}$ is bounded on N_σ . Moreover, one can view $(\hat{E}_\sigma)^{\dagger}$ as a dense defined unbounded operator on N_σ , in fact, it is the inverse operator of $\hat{E}_\sigma$.

For $\sigma \in R$, let j_σ be the operator $j_\sigma : N_\sigma \rightarrow N$, $(j_\sigma \psi)(t, x) = \psi(x) \delta(t - \sigma)$ and j_σ^* be its adjoint. If Λ is a closed subset of R^{n+1} we denote by N_Λ the subspace of N which comprises all distributions with support in Λ . The orthogonal projection of N in N_Λ will be denoted by e_Λ . Following [5] we have:

Proposition 1 *The operators j_σ are isometries and $j_\sigma^* j_\sigma = 1_{N_\sigma}$, $j_\sigma j_\sigma^* = e_\sigma$, where e_σ denotes the projector corresponding to the subset of R^{n+1} , $t = \sigma$.*

Then we define the operators:

$$U_{\sigma, \sigma'} : N_{\sigma'} \rightarrow N_\sigma, U_{\sigma, \sigma'} = j_\sigma^* \circ j_{\sigma'}. \quad (4)$$

We will derive in the following that $U_{\sigma, \sigma'}$ are propagators in the sense of [6]. This will follow from the Markoff property of the projectors e_σ .

Lemma 2 *Let A , B and C be closed subsets in R^{n+1} such that C separates A and B . Then $e_A \circ e_C \circ e_B = e_A \circ e_B$.*

Solution 3 *This is the consequence of the fact that E_m is the kernel of a local operator. The proof is identic with that of [5].*

The basics properties of $U_{\sigma,\sigma'}$ operators are stated in the following proposition.

Proposition 4 *The family of operators $U_{\sigma,\sigma'}$, $\sigma, \sigma' \in R$ has the following properties:*

- 1) $U_{\sigma,\sigma'} \circ U_{\sigma',\sigma''} = U_{\sigma,\sigma''}$
- 2) $U_{\sigma,\sigma} = 1_{N_\sigma}$
- 3) $\|U_{\sigma,\sigma'}\| \leq 1$.

Solution 5 1) *Using the Markoff property we have:*

$$e_\sigma \circ e_{\sigma'} \circ e_{\sigma''} = e_\sigma e_{\sigma''} \Leftrightarrow j_\sigma \circ j_\sigma^* \circ j_{\sigma'} \circ j_{\sigma'}^* \circ j_{\sigma''} \circ j_{\sigma''}^* = j_\sigma \circ j_\sigma^* \circ j_{\sigma''} \circ j_{\sigma''}^*. \quad (5)$$

By composition with $j_{\sigma''}$ at the right, we have

$$j_\sigma \circ (j_\sigma^* \circ j_{\sigma'} \circ j_{\sigma'}^* \circ j_{\sigma''} - j_\sigma^* \circ j_{\sigma''}) = 0. \quad (6)$$

From the definition of $U_{\sigma,\sigma'}$ and since j_σ are isometries, we conclude $U_{\sigma,\sigma'} U_{\sigma',\sigma''} = U_{\sigma,\sigma''}$.

2) *It follows from proposition 1.1 and definition of $U_{\sigma,\sigma'}$.*

3) *Because j_σ^* and j_σ are isometries, the property results immediately.*

III. EXPONENTIAL BOUNDS ON PROPAGATORS

To improve our estimates on the propagators $U_{\sigma,\sigma'}$ we need a supplementary condition on the metric g . We say that an application $Q : R^{n+1} \rightarrow M(n+1, n+1)$ has stable positivity if there exists $\varepsilon > 0$ such that for any application $\delta : R^{n+1} \rightarrow M(n+1, n+1)$ with $|\delta(x)^{ij}| \leq \varepsilon$ the matrices $Q(x) - \delta(x)$ are positive defined for any $x \in R^{n+1}$. The following result is a direct application of Agmon theory [2] of exponentially decay of solutions of elliptic second order operators.

Proposition 6 *If the metric g has stable positivity then for any $f \in N_{\sigma'}$:*

$$\int_{T_0}^{\infty} d\sigma \left\{ e^{\omega\sigma} \left\| \hat{E}_{\sigma}^{1/2} \circ U_{\sigma,\sigma'} f \right\|_{N_{\sigma}} \right\}^2 < \infty, \quad (7)$$

provided $\omega < \frac{m}{\sqrt{\sup g^{11}}}$.

Solution 7 *Starting from*

$$\begin{aligned} \langle u, U_{\sigma,\sigma'} f \rangle_{N_{\sigma}} &= \left\langle u, \hat{E}_{\sigma} \circ U_{\sigma,\sigma'} f \right\rangle_{L^2(R^n, d\mu_{\sigma})} \\ &= \int_{R^n} \bar{u}(x) \left[\int_{R^n} E_m(\sigma, x; \sigma', y) f(y) d\mu_{\sigma'}(y) \right] d\mu_{\sigma}(x) \end{aligned} \quad (8)$$

for $u \in N_{\sigma}$ and $f \in N_{\sigma'}$, it follows that $\varphi(\sigma, x) = \left(\hat{E}_{\sigma} \circ U_{\sigma,\sigma'} f \right)(x)$ is a solution of

$$(\Delta + m^2) \varphi(\sigma, x) = 0 \quad (9)$$

for $\sigma > \sigma'$. Let $\rho_m(\cdot; \cdot)$ denotes the distance corresponding to the metric $g_m = mg$. The metric g has stable positivity so, there is an $\varepsilon \in R_+$ such that $\rho_m(\sigma_0, x_0; \sigma, x) > \frac{\varepsilon}{m} |\sigma - \sigma_0|$. For $\Omega = \{(\sigma, x) : \sigma > T_0\}$, $T_0 \in R_+$ and for some positive λ :

$$\begin{aligned} &\int_{\Omega} |\varphi(\sigma, x)|^2 e^{-\lambda \rho_m(T_0, x_0; \sigma, x)} \sqrt{g(\sigma, x)} d\sigma d^n x \\ &= \int_{T_0}^{\infty} d\sigma \left\langle \hat{E}_{\sigma} \circ U_{\sigma,\sigma'} f, \hat{E}_{\sigma} \circ U_{\sigma,\sigma'} f \right\rangle_{L^2(R^n, d\mu_{\sigma})} e^{-\lambda \frac{\varepsilon}{m} (\sigma - T_0)} \\ &< ct. \int_{T_0}^{\infty} d\sigma \left\langle U_{\sigma,\sigma'} f, \hat{E}_{\sigma} \circ U_{\sigma,\sigma'} f \right\rangle_{L^2(R^n, d\mu_{\sigma})} e^{-\lambda \frac{\varepsilon}{m} (\sigma - T_0)} \\ &= ct. \int_{T_0}^{\infty} d\sigma \|U_{\sigma,\sigma'} f\|_{N_{\sigma}}^2 e^{-\lambda \frac{\varepsilon}{m} (\sigma - T_0)} < \infty. \end{aligned} \quad (10)$$

So we are in the conditions of the main theorem of [2]. It follows that:

$$\begin{aligned} &\int_{\Omega} d\sigma d^n x \sqrt{g(\sigma, x)} |\varphi(\sigma, x)|^2 (m^2 - g(\nabla h(\sigma, x), \nabla h(\sigma, x))) e^{2h(\sigma, x)} \\ &\leq \frac{2(1+2d)}{d^2} m^2 \int_{\Omega \setminus \Omega_d} |\varphi(\sigma, x)|^2 e^{2h(\sigma, x)} \sqrt{g(\sigma, x)} dx, \end{aligned} \quad (11)$$

where d is a positive number and $\Omega_d = \{(\sigma, x) \in \Omega : \rho_m((\sigma, x), \{\infty\}) > d\}$. Here

$$\rho_m((\sigma, x), \{\infty\}) = \sup \{ \rho_m((\sigma, x), \Omega \setminus K) : K \text{ is a compact subset of } \Omega \}. \quad (12)$$

The function h is any function which satisfies the condition $g(\nabla h(\sigma, x), \nabla h(\sigma, x)) < m^2$.

We choose $h(\sigma, x) = \omega\sigma$ with $\omega < \frac{m}{\sqrt{\sup g^{11}}}$. The above inequality becomes

$$\begin{aligned}
& \int_{\Omega} d\sigma d^n x \sqrt{g(\sigma, x)} |\varphi(\sigma, x)|^2 e^{2\omega\sigma} \\
& < \frac{2(1+2d)}{d^2} \frac{m^2}{m^2 - \omega^2} \int_{\Omega \setminus \Omega_d} d\sigma dx \sqrt{g(\sigma, x)} |\varphi(\sigma, x)|^2 e^{2\omega\sigma}.
\end{aligned} \tag{13}$$

If for any point $(\sigma, x) \in \Omega$ there is a geodesic which starts in (σ, x) and ends in the hyperplane $\sigma = T_0$ then $\Omega \setminus \Omega_d \subset \{(\tau, x) : 0 < \sigma \leq T\}$ with T sufficiently large but finite. In conclusion

$$\begin{aligned}
& \int_{\Omega} d\sigma d^n x \sqrt{g(\sigma, x)} |\varphi(\tau, x)|^2 e^{2\omega\sigma} \\
& = \int_{T_0}^{\infty} d\sigma e^{2\omega\sigma} \left\langle \hat{E}_{\sigma} \circ U_{\sigma, \sigma'} f, \hat{E}_{\sigma} \circ U_{\sigma, \sigma'} f \right\rangle_{L^2(R^n, \mu_{\sigma})} < \infty,
\end{aligned} \tag{14}$$

or

$$\int_{T_0}^{\infty} d\sigma e^{2\omega\sigma} \left\langle \hat{E}_{\sigma} \circ U_{\sigma, \sigma'} f, \hat{E}_{\sigma} \circ U_{\sigma, \sigma'} f \right\rangle_{L^2(R^n, \mu_{\sigma})} < \infty, \tag{15}$$

which implies

$$\int_{T_0}^{\infty} d\sigma \left\{ e^{\omega\sigma} \left\| \hat{E}_{\sigma}^{1/2} \circ U_{\sigma, \sigma'} f \right\|_{N_{\sigma}} \right\}^2 < \infty. \tag{16}$$

IV. THE STATIONARY CASE

We consider in this section that there is a coordinate system such that the metric g is independent of first coordinate. In this case, the spaces N_{σ} and the operators $\hat{E}_{\sigma}^{1/2}$ are identically and will be denoted by N_0 and $\hat{E}_0^{1/2}$ respectively. Thus, the operators $U_{\sigma, \sigma'}$ are defined on the same Hilbert space and depend only on the difference $\sigma - \sigma' : U_{\sigma, \sigma'} = U_{\sigma - \sigma'}$. The family of operators $\{U_{\tau}\}_{\tau \in R_+}$ forms a semigroup. Using the results about existence and properties of the generators of semigroups [7], we can obtain bounds directly on the transfer matrix U_{τ} .

Proposition 8 *The semigroup $\{U_{\tau}\}_{\tau \in R_+}$ is exponentially bounded: $\|U_{\tau}\|_{N_0} < e^{-\tau\omega}$ provided $\omega < \frac{m}{\sqrt{\sup g^{11}}}$.*

Solution 9 *Because we have found estimates on $\hat{E}_0^{1/2} \circ U_{\tau}$, we will consider the operators $\tilde{U}_{\tau} = \hat{E}_0^{1/2} \circ U_{\tau} \circ (\hat{E}_0^{1/2})^{\dagger}$, well defined on $L^2(R^n, d\mu_0)$. Using the fact that $L^2(R^n, d\mu_0)$ is*

dense in N_0 we can extend these operators by continuity on the space N_0 . In this way we have build the semigroup $\left\{\tilde{U}_\tau\right\}_{\tau \in R_+}$ which satisfies the estimates of the precedent section:

$$\int_{T_0}^{\infty} d\tau \left\{ e^{\omega\tau} \left\| \tilde{U}_\tau \right\|_{N_0} \right\}^2 < \infty, \quad (17)$$

for some $T_0 > 0$. So $\left\{\tilde{U}_\tau\right\}_{\tau \in R_+}$ is exponentially bounded and in consequence [7], if $\tilde{K}$ is its generator ($\tilde{U}_\tau = e^{-\tau\tilde{K}}$) the resolvent set of $\tilde{K}$ satisfies:

$$\{z \in C \mid \operatorname{Re} z \in (-\infty, \omega)\} \subset \rho(\tilde{K}). \quad (18)$$

If K is the generator of $\{U_\tau\}_{\tau \in R_+}$ then, on $\mathcal{D}(K)$ we have:

$$K = \left(\hat{E}_0^{1/2}\right)^\dagger \circ \tilde{K} \circ \hat{E}_0^{1/2} \quad (19)$$

by using the reciprocal formula

$$U_\tau = \left(\hat{E}_0^{1/2}\right)^\dagger \circ \tilde{U}_\tau \circ \hat{E}_0^{1/2}, \quad (20)$$

valid on N_0 . If the operator

$$\left(\hat{E}_0^{1/2}\right)^\dagger \circ \left(\tilde{K} - z\right)^{-1} \circ \hat{E}_0^{1/2} \quad (21)$$

is well defined, even on a dense subset of N_0 , then $K - z$ is inversable. From 20 it follows that, if $\left(\tilde{K} - z\right)^{-1}$ exists, then:

$$\left(\tilde{K} - z\right)^{-1} \left(L^2(R^n, d\mu_0)\right) \subset L^2(R^n, d\mu_0), \quad (22)$$

and in consequence $\left(\hat{E}_0^{1/2}\right)^\dagger \circ \left(\tilde{K} - z\right)^{-1} \circ \hat{E}_0^{1/2}$ is well defined on the entire N_0 . Will follow that $\rho(\tilde{K}) \subset \rho(K)$ and this ends the proof.

If the metric is symmetric at transformation $x^1 \rightarrow -x^1$, the transfer matrix generator is self-adjoint and it can be considered as the Hamiltonian of the scalar field.

V. APPLICATION

Our application is for the Euclidean case. The results concerning decoupling of different regions in quantum Euclidean fields are based primarily on estimates of $\|e_{\Lambda_1}e_{\Lambda_2}\|_N$, where Λ_1, Λ_2 are two disjoint regions. Let us consider the two dimensional case. The most difficult case is when Λ_1, Λ_2 are not convex and there is no possibility of drawing a straight line between the two subsets. We can sharpen the existent estimates [5] for these cases by using the previous results. The idea is to make a change of coordinates such that for the new coordinates, lines like $\sigma = ct$. separate the two sets and they are as closed as possible to the boundaries of Λ_1, Λ_2 . Then we can use the exponential bounds of the previous section to evaluate $\|e_{\Lambda_1}e_{\Lambda_2}\|_N$. More precisely:

Proposition 10 *Let Λ_1, Λ_2 two regions in R^2 such that the construction of the coordinates 24 to be possible (after a rotation if necessary). Then*

$$\|e_{\Lambda_1} \circ e_{\Lambda_2}\|_N \leq e^{-m|\beta-\alpha|\min|\cos\theta|}, \quad (23)$$

where θ and $|\beta - \alpha|$ will be defined during the proof.

Solution 11 *Let (t, x) denotes the original coordinates in which the metric is diagonal. Let $\gamma : R \rightarrow R^2$ be a curve which separates Λ_1, Λ_2 and $\gamma(0) = (t = 0, x = 0)$. We define a new coordinate system (σ, ξ) by*

$$\begin{cases} t(\sigma, \xi) = \sigma + \gamma^1(\xi) \\ x(\sigma, \xi) = \gamma^2(\xi) \end{cases} \quad (24)$$

In the new coordinates, the metric is

$$g'(\sigma, \xi) = \begin{pmatrix} 1 & \frac{d\gamma^1}{d\xi} \\ \frac{d\gamma^1}{d\xi} & \left(\frac{d\gamma^1}{d\xi}\right)^2 + \left(\frac{d\gamma^2}{d\xi}\right)^2 \end{pmatrix} \quad (25)$$

so we are in the conditions of the last section. Using the Markoff property,

$$\|e_{\Lambda_1} \circ e_{\Lambda_2}\|_N = \|e_{\Lambda_1} \circ e_\alpha \circ e_\beta \circ e_{\Lambda_2}\|_N \leq \|e_\alpha \circ e_\beta\|_N, \quad (26)$$

where the lines $\sigma = \alpha$, $\sigma = \beta$ separate Λ_1 and Λ_2 exactly in the order they appear in the above relation (in the sense that $\sigma = \alpha$ separates Λ_1 by $\sigma = \beta$ etc.). Further

$$\left\|j_\alpha \circ j_\alpha^\dagger \circ j_\beta \circ j_\beta^\dagger\right\|_N = \left\|j_\alpha \circ U_{\alpha-\beta} \circ j_\beta^\dagger\right\|_N = \|U_{\alpha-\beta}\|_{N_0}. \quad (27)$$

The element $(g')^{11}$ is given by $(g')^{11} = \frac{1}{\cos^2 \theta}$, where θ is the angle between the tangent to the curve γ and the x axis. Using the bounds of the last section we have

$$\|e_{\Lambda_1} \circ e_{\Lambda_2}\|_N \leq e^{-m|\beta-\alpha| \min|\cos \theta|}. \quad (28)$$

Performing first a rotation, one can choose the best values for $|\beta - \alpha|$ and $\min|\cos \theta|$.

VI. CONCLUSIONS

Our primary goal was to define the transfer matrix for scalar fields on curved spaces and to investigate the basic spectral properties of its generator. Even though the generator is not self-adjoint in the general case, this approach allows us to investigate this problem by using at least two new tools besides the methods of Green functions. One is the perturbations of hypercontractive semigroups [8] and the other is the adiabatic theorem.

Now it is straightforward to quantize the field by defining the Markoff field over the space N . For the stationary, symmetric at time reflection case (static), we think that one has now all elements to construct the physical field (for example that proposed in [4]) by following Nelson reconstruction method and holomorphic continuation of the transfer matrix. Note that, according to results of [6], the holomorphic continuation of the transfer matrix to real time is still possible, in the stationary case without symmetry at time reflection, as long the spectrum of the generator belongs to the real axis. Of course, one has to check that the results of [9] (sistemized in [5]), which are the core of the reconstruction theorem, are still valid. For the general case, we think that the adiabatic theorem, especially the adiabatic reduction theory [10], may play an important role in defining the physical quantum field by following Nelson's approach.

REFERENCES

- [1] Nelson E., J. Func. Anal. **12**, 97 (1973)
- [2] Agmon S., *Lectures on exponential decay of solutions of second-order elliptic equations*, Princeton: Princeton Univ. Press (1982)
- [3] Guerra F., Rosen L., Simon B., Ann. Math. **101**, 111 (1975)
- [4] Dimock J., J. Math. Phys. **20**, 2549 (1979)
- [5] Simon B., *The $P(\phi)_2$ Euclidean (Quantum) Field Theory*, Princeton: Princeton University Press (1974)
- [6] Reed M., Simon B., *Methods of Modern Mathematical Physics*, Vol. 2, New York: Academic Press (1975)
- [7] Neerven J., *The asymptotic behavior of semigroups of linear operators*, Basel, Boston: Birkhauser (1996)
- [8] Simon B., Hoegh-Krohn R., J. Func. Anal. **9**, 121 (1972)
- [9] Nelson E., J. Func. Anal. **11**, 211 (1972)
- [10] Nenciu G., Commun. Math. Phys. **152**, 479 (1993)